



On the Evolution of a Sub-C Class Flare: A Showcase for the Capabilities of the Revamped Catania Solar Telescope

Paolo Romano¹ · Salvo L. Guglielmino¹ · Pierfrancesco Costa¹ · Mariachiara Falco¹ · Salvatore Buttaccio¹ · Alessandro Costa¹ · Eugenio Martinetti¹ · Giovanni Occhipinti¹ · Daniele Spadaro¹ · Rita Ventura¹ · Giuseppe E. Capuano² · Francesca Zuccarello²

Received: 6 October 2021 / Accepted: 30 November 2021
© The Author(s), under exclusive licence to Springer Nature B.V. 2022

Abstract

Solar flares are occasionally responsible for severe space-weather events, which can affect space-borne and ground-based infrastructures, endangering anthropic technological activities and even human health and safety. Thus, an essential activity in the framework of space-weather monitoring is devoted to the observation of the activity level on the Sun. In this context, the acquisition system of the Catania Solar Telescope has been recently upgraded in order to improve its contribution to the European Space Agency (ESA) – Space Weather Service Network through the ESA Portal, which represents the main asset for space weather in Europe. Here, we describe the hardware and software upgrades of the Catania Solar Telescope and the main data products provided by this facility, which include full-disk images of the photosphere and chromosphere, together with a detailed characterization of sunspot groups. As a showcase of the observational capabilities of the revamped Catania Solar Telescope, we report the analysis of a B5.4 class flare that occurred on 7 December 2020, simultaneously observed by the Interface Region Imaging Spectrograph and the Solar Dynamics Observatory satellites.

Keywords Instrumentation and data management · Flares · Dynamics

1. Introduction

Solar eruptions are the most prominent manifestations of the magnetic activity of the Sun, involving the entire heliosphere, with a potential impact on the Earth (e.g. Patsourakos et al., 2016; Piersanti et al., 2017). Their effects on the near-Earth environment and anthropic

✉ P. Romano
paolo.romano@inaf.it

S.L. Guglielmino
salvatore.guglielmino@inaf.it

¹ INAF - Osservatorio Astrofisico di Catania, Via S. Sofia 78, 95123 Catania, Italy

² Dipartimento di Fisica e Astronomia “Ettore Majorana” – Sezione Astrofisica, Università degli Studi di Catania, Via S. Sofia 78, 95123 Catania, Italy

activities are increasingly involving socioeconomical interests, taking into account that solar eruptions may provoke severe damage to technological systems. Therefore, many recent efforts of the scientific community have been addressed to forecast eruptive events with enough advance to prevent or mitigate their impact on human activities (Schwenn, 2006).

Eruptive phenomena are usually associated with flares, which are sudden energy-release events that emit radiation across the entire electromagnetic spectrum, as well as accelerated particles (see Benz, 2017, for a review). Like eruptions, flares occur because of the release of free magnetic energy, which is converted into heat and kinetic energy through magnetic reconnection (Kopp and Pneuman, 1976; Moore et al., 2001). This leads to the observed brightness enhancements that can be easily detected also at visible wavelengths, such as the $H\alpha$ line, centered at $\lambda = 656.28$ nm, which is one of the most popular lines for studying the solar chromosphere (see Leenaarts, Carlsson, and Rouppe van der Voort, 2012, and references therein). The flaring chromosphere is characterized by strong $H\alpha$ emission, related to the impact of energetic particles accelerated at the reconnection site colliding in the chromospheric plasma. $H\alpha$ emission often appears as two bright ribbons, separating from each other (see, e.g., Fletcher et al., 2011). However, considering the three-dimensional nature of magnetic reconnection (Aulanier, Janvier, and Schmieder, 2012; Aulanier et al., 2013), the presence of a more complex topology in the reconnection site may result in flares with extra ribbons or even circular ribbons (e.g. Masson et al., 2009; Guglielmino et al., 2016; Romano et al., 2017).

Although the trigger of flares is thought to be located in the upper layers of the solar atmosphere, with the coronal magnetic field playing a leading role, most of the flare-forecasting methods are based on photospheric observations of the active regions (ARs) where flares occur. Indeed, the magnetic-field configuration suitable for the occurrence of eruptive events is mainly driven by the photospheric evolution of the ARs, determined by the emergence of new magnetic flux from the convection zone into the solar atmosphere and by the rearrangement of the coronal field due to the horizontal photospheric displacements of the field-line footpoints (e.g. Romano and Zuccarello, 2007; Romano et al., 2015, 2018; Romano, Elmhamdi, and Kordi, 2019), leading to magnetic helicity changes (e.g. Zuccarello et al., 2021). Thus, many approaches to flare forecasting produced in the last decades rely on the determination of certain photospheric parameters, such as the total unsigned magnetic flux together with the length of the magnetic polarity-inversion line (PIL) characterized by a strong line-of-sight field gradient and the total magnetic-energy dissipation (Yuan et al., 2010), or with the free magnetic energy obtained from the line-of-sight field gradient along the PILs (Falconer, Moore, and Gary, 2008). More recently, Korsós et al. (2015) proposed the introduction of the weighted horizontal magnetic gradient (WG_M) to forecast flares based on the magnetic gradient among all spots within an appropriately defined region close to the PIL (see also Korsós and Erdélyi, 2016; Korsós, Yang, and Erdélyi, 2019).

Other forecasting methods are performed on a statistical basis. Bloomfield et al. (2012) proposed a method that uses the McIntosh group classification of sunspot groups (SGs) observed in the photosphere and assumes that flares are Poisson-distributed processes; these authors showed that Poisson probabilities perform comparably better than other more complex prediction methods. The extensive comparison among several flare-forecasting methods performed by Barnes et al. (2016) suggested that it may be possible to obtain the best prediction by combining a method that characterizes an AR by one or more parameters, and uses a statistical technique. In this perspective, Falco, Costa, and Romano (2019) developed a method that is based mainly on the Zurich classification of the SGs observed at the photospheric level and assumes Poisson statistics for the flare occurrence. This method is able to provide an estimation of the capability to host flares of a specified energy range for an AR characterized by a particular configuration, size, and fragmentation.

In this article, we present the capabilities of the revamped Catania Solar Telescope. This facility, built in the 1960s, is now able to provide almost simultaneous full-disk observations of the chromosphere and photosphere in the $H\alpha$ line and in the nearby continuum with high cadence, down to 1 s, with an angular resolution of $2''$. Such observations are an effective tool for a synoptic monitoring of the flaring activity of the chromosphere, as well as for characterizing several photospheric parameters of the ARs where flares occur.

The article is organized as follows. In Section 2 we describe the upgrade of the Catania Solar Telescope. Section 3 reports on the observation of a B5-class flare, simultaneously observed by the Catania Solar Telescope and by the Solar Dynamics Observatory (SDO) and the Interface Region Imaging Spectrograph (IRIS) satellites, as a showcase for illustrating the capabilities of the revamped facility. In Section 4 we draw our conclusions.

2. The Catania Solar Telescope

The observations of the solar photosphere and chromosphere are carried out at the Istituto Nazionale di Astrofisica (National Institute for Astrophysics) – Catania Astrophysical Observatory (INAF-OACT), in collaboration with the University of Catania, by means of a telescope equipped with two refractors with diameters of 150 mm and focal lengths of 2230 mm and 2300 mm, respectively. The first one is used to make drawings of sunspot groups and pores from visual observations, the second feeds a Zeiss Lyot filter (bandwidth of 0.025 or 0.050 nm, tunable filter range ± 0.1 nm), which is used to take digital full-disk images of the photosphere in the continuum of the $H\alpha$ line at 656.78 nm and of the chromosphere in the center of the $H\alpha$ line at 656.28 nm.

On a daily basis, when the weather conditions allow, a drawing of the projected Sun is performed in order to determine some properties of the sunspot groups visible in the photosphere for each of them: heliographic latitude and longitude of the barycenter, number of sunspots and pores, projected area in tens of millionths of the solar hemisphere, type of penumbra of the main sunspot of the group, relative importance between the leading spot and density of the sunspot population (see Ternullo et al., 2006), and group type according to the Zurich classification (Falco, Costa, and Romano, 2019). Given the characteristics of the instrument and average seeing conditions, the number of pores that can be detected by visual inspection of the projected photospheric image becomes greater than that retrieved by the digitized images, thanks to the better resolution of the former option. The photospheric data acquired at the OACT are distributed to international Solar Data Centers, like the Solar Influences Data Center (SIDC) in Brussels and the World Data Center for the Sunspot Index (NOAA, Boulder).

When the weather conditions permit, digital images are also acquired in the center of the $H\alpha$ line and in the nearby continuum by the Lyot filter. The OACT contributes with its $H\alpha$ images to the Global High Resolution $H\alpha$ Network. The digital images of the Sun are also provided, on request, to observers carrying out observational campaigns using high-resolution solar telescopes.

The sunspot group characterization, the photospheric, and the chromospheric data are published in near real time on the portal of the ESA Space Situational Awareness Programme (<https://swe.ssa.esa.int/solar-weather>).

2.1. Hardware Upgrade

From October 2012 to September 2020 a 3056×3056 Kodak KAF-9000 CCD array was in operation for digital image acquisition, with a dynamical range of 16 bit, a pixel size of

12 μm , and a time resolution of about 150 s. A noise of 12 e^- RMS and a dark current $< 1.5 e^- \text{pixel}^{-1} \text{s}^{-1}$ characterized that detector.

In order to improve the quality of the images and their time resolution, in September 2020 a new detector was installed at the focal plane of the telescope. A back-illuminated sCMOS camera with a sensor of 2048×2048 pixels, a pixel size of 11 μm , and a quantum efficiency of 95% revamped the capabilities of the telescope. In particular, the new fast acquisition capability of this device allows us to follow the dynamic solar processes occurring in the chromosphere in near real time. We changed the default acquisition rate in the center of the $H\alpha$ line from one image per 10 minutes to one image per minute. Moreover, with a maximum frame rate of 24 and 48 fps at 12 and 16 bit, respectively, new applications and new restoring techniques can be used to improve the image quality (e.g. using the lucky imaging technique; see Law, Baldwin, and Mackay, 2006; Mackay, 2013).

We also modified the system by introducing an electronic shutter. The use of an on-sensor rolling shutter overcomes the need for mechanical shutters. This avoids the exposure gradient effects associated with those induced by, e.g., an iris shutter, thus providing much better accuracy for photometry. Moreover, the rolling shutter allows using shorter time exposures and adapting them on the basis of the sky conditions, as well as on the level of solar activity.

The above-mentioned upgrades allowed us to reduce the exposure time from 0.2 s, used with the old setup, down to 0.004 s currently operated.

An example of an image taken during good seeing conditions can be seen in Figure 1.

2.2. Software Upgrade

In addition, an upgrade of the pipeline for the acquisition and handling of the data has been carried out. This aspect allowed us to improve the quality of the raw and corrected images. By the new acquisition system, raw images are first transferred to the machine dedicated to the preliminary analysis and data reduction, afterwards they are moved to the storage server (Figure 2).

The raw images (red boxes in Figure 2) acquired by the telescope are of three types: $H\alpha$ images taken in order to obtain a flat field image by the KLL method (Kuhn, Lin, and Lorz, 1991), $H\alpha$, and continuum scientific images for observations of the chromosphere and photosphere, respectively.

An $H\alpha$ sequence for a flat-field purpose is taken every day in about one minute at the beginning of the observing run. The sequence is formed by nine images: one image taken with the center of the solar disk close to the center of the detector, four images with the solar disk almost tangent to each of the four sides of the detector, and four images with the solar disk near each of the four corners of the detector. These images are obtained by changing the pointing of the telescope. Assuming that the Sun is a constant light source in one minute, the KLL algorithm uses each pair of pixels illuminated by the same part of the Sun in successive displaced pictures to measure the ratio of the gain of those pixels. By an iteration process it is possible to reconstruct the pixel-to-pixel nonuniformity in the gain of the whole detector (e.g. Li et al., 2021). Therefore, to determine the center and radius of the Sun in the recorded images and to derive the relative displacements among the Sun images, a fitting of the solar limb is performed by an IDL routine. After the application of the KLL algorithm the output flat-field image, together with the input sequence, are stored in our archive.

By default, every day when weather conditions permit, from 8:00 CET to 13:30 CET, images in the line center of $H\alpha$ are acquired at a cadence of one image per minute. However, a higher cadence (up to one image per second) is taken when a flare is observed in the chromosphere. Currently, the switch to the faster acquisition procedure is performed manually by the observer, however, an automatic detection algorithm for flare occurrence based




Figure 1 Example of full-disk $H\alpha$ image taken by the Catania Solar Telescope during good seeing conditions and corrected by the flat field. Here, and in the following figures, solar north is at the top, and west is to the right.

on the near real-time analysis of the $H\alpha$ signal above the solar disk is under development and will be released in the near future (see the violet box in Figure 2).

After the limb fit and the determination of the center and radius of the solar disk, a quality-check procedure is applied to the image. A quality index from 1 to 3 (1=good, 3=bad) is assigned to each image based on the reliability of the solar-radius estimation and on the presence of some anomalies in the measured signal. For instance, when a cloud passes in front of the solar disk a quality index equal to 2 or 3 is assigned to the acquired image depending on the significance of the anomalous decrease of the signal. Regardless of the assigned quality index, further keywords are added to the preliminary header, which is initially generated by the acquisition program. These keywords are in agreement with the metadata definition for Solar Orbiter science data (De Groof, Walsh, and Williams, 2019). The output FITS format data are stored in the archive while the database is updated, accordingly. Indeed, a Ruby script is designed to ease the process of populating the database with the data records. The database containing the records of the images and metadata has been created in MySQL. The same steps are applied also to the continuum images.




Figure 2 Flow chart describing the new acquisition software operating at the INAF-Catania Solar Telescope.

Only images of quality class 1 (good) are processed further and additional jpeg images are produced. For the data taken in the center of the H α line, the contrast of the limb structures (e.g. solar prominences) is enhanced in order to increase the visibility of their morphology. Then, both chromospheric and photospheric images are corrected by the flat field over the solar disk and an additional contrast enhancement is performed. In the JPEG2000 images (2048 \times 2048 pixels), the solar disk is centered and institutional logos and a time stamp are added on top. The pipeline produces two versions of JPEG2000 files: with and without the heliographic grid plotted over the solar disk (the former in red scale for the H α images).

Only images of quality class 1 are available in the ESA Space Weather Service Network, while all images are available in our database (<http://ssa.oact.inaf.it/oact/index.html>).

3. Observation of a B5-Class Solar Flare

Solar flares involve different layers of the solar atmosphere and cover a wide range of energy levels and sizes. This allows us to investigate the physical processes related to the release of energy at different scales. However, the study of small flares has some advantages compared to major events, as flares characterized by a weaker emission do not saturate the digital images and allow analysis of the topology of the involved magnetic systems, like the flare-loop configuration or the shape of ribbons, in more detail. Moreover, small flares can be observed entirely within the field of view of high-resolution instruments. However, their unpredictable character makes it difficult to acquire a good quality dataset.

As a showcase of the observational capabilities of the revamped Catania Solar Telescope, we studied a small flare of B5.4 GOES class that occurred in active region (AR)

NOAA 12790 (hereafter, AR 12790) on 7 December 2020 with a peak at 09:55 UT. We used data acquired by the Catania Solar Telescope and by satellite instruments, such as the Interface Region Imaging Spectrograph (IRIS: De Pontieu et al., 2014), and the Helioseismic and Magnetic Imager (HMI: Scherrer et al., 2012) and Atmospheric Imaging Assembly (AIA: Lemen et al., 2012) on board the Solar Dynamics Observatory (SDO: Pesnell, Thompson, and Chamberlin, 2012).

3.1. Dataset Description

We used a sequence of images taken in the center of the $H\alpha$ line at 656.28 nm by the Catania Solar Telescope on 7 December 2020 from 9:30 UT to 10:30 UT. These images with a time cadence of 1 min were characterized by a spatial resolution of about $2''$ due to the average seeing conditions during the acquisition of that image sequence. Only four images have been neglected in our analysis for their low quality due to clouds in front of the solar disk between 10:01 UT and 10:05 UT.

We also analyzed a simultaneous observing sequence acquired by the IRIS satellite between 09:25:47 UT and 10:14:52 UT on 7 December. The sequence consisted of a single large dense 320-step raster scan (OBS3610108077). The sequence had a $0.33''$ step size and a 9.2 s step cadence, with a pixel size of $0.35''$ along the y -direction (spatial binned data), covering a field of view (FOV) of $112'' \times 175''$. Simultaneously, slit-jaw images (SJIs) were acquired in the 1400, 1330, and 2796 Å passbands, corresponding to the Si IV 1402 Å, C II 1335 Å, and Mg II k lines, respectively. These SJIs have a cadence of 37 s for consecutive frames in each passband and cover a FOV of $167'' \times 175''$.

To determine the magnetic context of the flaring active region, we used photospheric observations from the SDO satellite consisting of full-disk continuum filtergrams and line-of-sight (LOS) magnetograms taken by HMI along the Fe I 6173 Å line, with a spatial resolution of $1''$. Furthermore, we took advantage of the coronal images acquired by AIA using the EUV filters centered at 131 Å, 193 Å, 171 Å, and 304 Å and UV filters at 1600 Å and 1700 Å. These EUV/UV data have an image spatial scale of about $0.6''$ per pixel and a cadence of 12 s.

The alignment between the $H\alpha$ images acquired with the Catania Solar Telescope and those acquired by the IRIS and SDO satellites was obtained by applying the IDL SolarSoft mapping routines and crosscorrelation techniques with respect to the cospatial sub-FOV between the different instruments (see Figure 3). We used the leading spot of AR 12790 within the sub-FOV as a fiducial point, taking into account the pixel scale of the different instruments. The accuracy of the alignment is $\pm 1''$, being comparable to the spatial resolution of SDO/HMI data.

3.2. Results

The analyzed flaring event (i.e. SOL2020-12-07T09:55) was part of a sequence of homologous precursor flares (see Romano et al., 2015) occurring in AR 12790, culminating in a C7.4 flare with a peak at 16:32 UT on the same day. A list of these events is provided in Table 1. The X-ray flux measured by the GOES-16 satellite between 6 December at 21:00 UT and 7 December at 21:00 UT is shown in Figure 4 (left panel). A sequence of images acquired with the AIA UV filter at 1600 Å that are relevant to the precursor flares, including the SOL2020-12-07T09:55 event, together with the stronger C7-class flare, is shown in Figure 4 (right panels). Comparing the location of the flare to the magnetic configuration of AR 12790 reported in Figure 3, we can see that the flaring emission during the precursor




Figure 3 Context images for the Catania Solar Telescope observations relevant to AR 12790. Left panel: Full-disk $H\alpha$ image acquired with the Catania Solar Telescope during the B5.4 flare. The solid box frames the FOV analyzed in the right panels. Right panels: Maps of SDO/HMI continuum intensity (top) and magnetogram (bottom) for the FOV indicated with a solid box in the full-disk $H\alpha$ image. The dashed lines enclose the sub-FOV further investigated in the article and shown in the following figures.

Table 1 Flaring events occurring in AR 12790, belonging to the sequence of homologous precursor flares described in the main text. The event observed by the Catania Solar Telescope is indicated using bold letters.

Event	Date	Hour peak	Flare class
1	06-Dec-20	23:42 UT	B8.3
2	07-Dec-20	09:55 UT	B5.4
3	07-Dec-20	14:38 UT	B6.0
4	07-Dec-20	16:32 UT	C7.4

events takes place along the PIL of AR 12790, with a compact ribbon aligned along the north–south direction. As we can infer from the comparison between the $H\alpha$ image and SDO/HMI magnetogram, a filament characterized by an elongated S-shape is located along the PIL. The northern and southern ends of this filament correspond to the eastern side (towards the negative x -axis) of the preceding sunspot of the AR and the southern magnetic-field concentrations of the AR, respectively (see Figure 3). The location of the flare ribbons suggests that only a portion of the filament is involved in the recurrent flares, which are not able to completely destabilize the main magnetic-field configuration supporting the filament.

Figure 5 displays simultaneous multiwavelength observations acquired at the peak of the SOL2020-12-07T09:55 event, during the time interval covered by the Catania Solar Telescope observations. A compact ribbon is observed at decreasing temperature formation heights, according to the response of AIA EUV and UV filters, taking into account the emission contributions due to flares (O'Dwyer et al., 2010). At 304 \AA a bright patch protruding from the ribbon is observed near the preceding sunspot of AR 12790. This emitting structure is also faintly visible in the $H\alpha$ map (see the right panel of Figure 5), and it is clearly detected in the simultaneous IRIS SJIs at 1400 \AA (not shown here). Probably, this structure, which is visible mainly at a chromospheric level, can be interpreted as the emission produced in the northern footpoint of the filament by the electron beams, characterized by particular high




Figure 4 Left panel: Plot of the GOES-16 X-ray flux in the 1–8 Å passband from 6 December 2020 at 21:00 UT until 7 December 2020 at 21:00 UT. The yellow-shaded area indicates the time interval of the Catania Solar Telescope observations. Events listed in Table 1 are indicated with arrows and numbers. Right panels: SDO/AIA 1600 Å filtergrams related to the peak of the flares belonging to the sequence listed in Table 1.

energy and reaching the lower layers of the solar atmosphere. For this reason, probably, this bright region is not visible in the corona by AIA EUV filters.

The sub-FOV indicated with a dashed box in the H α map of Figure 5 has been used to compute the light curves of the SOL2020-12-07T09:55 flare in different wavelengths. Figure 6 shows the light curve deduced from the H α sequence (black symbols), as well as the light curve relevant to the IRIS SJIs used in this work (1400, 1330, and 2976 Å, colored symbols). The H α light curve has been computed by dividing the average value of the brightness in the sub-FOV by the brightness value at the disk center, in order to eliminate global observational effects, such as the presence of clouds. Therefore, a value larger than 1 at the beginning of the light curve indicates that we were observing a facula region. The background value for IRIS has been set by considering the average value in the sub-FOV at the beginning of SJI observations. For ease of comparison, we also plot in the same graph the GOES-16 X-ray flux in the 1–8 Å (blue line). Despite the small intensity of the flare (B5.4 GOES class), we note that the sensitivity of the Catania Solar Telescope is able to detect the increase of the brightness in the center of the H α line and to highlight the almost contemporary occurrence of the light curve peak in comparison to the IRIS wavelengths.

In Figure 7 we illustrate the time delay between the intensity peak occurring in the light curves relative to the SDO/AIA 304 Å channel for the compact ribbon and the remote ribbon of the flare, respectively. This time delay is 120 s, indicating a significant travel time for the propagation of the disturbance in comparison to a classical spine configuration (Masson et al., 2012).

Finally, in Figure 8 we report the observations relative to the IRIS SJIs for the restricted sub-FOV indicated in Figure 5. These high-resolution images allow us to obtain a better insight into the fine structure of the compact ribbon of the SOL2020-12-07T09:55 flare.




Figure 5 Simultaneous multiwavelength observations during the peak of the SOL2020-12-07T09:55 flare. Left panels: Maps derived from SDO/AIA observations corresponding to filters with decreasing temperature formation, from EUV 131 Å down to UV 1700 Å. Right panel: H α observation acquired with the Catania Solar Telescope. The dashed box in the H α map indicates the sub-FOV used for computing the light curve shown in Figure 6 and for studying the sequence of IRIS observations shown in Figure 8.




Figure 6 Light curves computed from simultaneous multiwavelength observations during the SOL2020-12-07T09:55 flare: H α data acquired with the Catania Solar Telescope (black symbols) and IRIS SJIs at 1400, 1330, and 2796 Å (see the inset), together with the GOES-16 X-ray flux in the 1–8 Å. All the curves are referred to the sub-FOV indicated with a dashed box in the H α map in Figure 5.

Indeed, the compact ribbon appears to consist of threads, especially in its northern part. At the beginning of the sequence (09:49 UT) in the Mg II k (2796 Å) images, the dark




Figure 7 Light curves computed from AIA 304 Å filtergrams, relevant to the boxes indicated in the inset. Note that the solid box, framing the compact ribbon, encloses the same sub-FOV indicated with a dashed box in the H α map as in Figure 5. The dashed box encloses the remote ribbon. For comparison, we include the GOES-16 X-ray flux in the 1–8 Å. The vertical continuum and dashed lines indicate the times corresponding to the emission peak of the main and northern flare ribbons, respectively.

filament shows its twisted fine structure, while its counterpart is very thin and bright in the 1400 and 1330 Å passbands. After the peak, when all the IRIS SJIs are saturated and the compact ribbon is observed, it is possible to detect a bifurcation of the filament, especially in the C II (1330 Å) and Mg II k images. This behavior is typical of the homologous flares characterized by the reformation of the involved filament after each event by the splitting of a single flux rope during the eruption (e.g. Gibson and Fan, 2006) or by the eruption of the upper part of a double-decker system (e.g. Kliem et al., 2014).

4. Discussion and Conclusions

In this article we have analyzed observations of a GOES B5-class flare occurring in a moderately complex active region and producing a compact ribbon at the chromospheric level and a farther remote ribbon. This event provides a contribution to the few observations of such multiple-ribbon events reported in the literature, which can be interpreted in the light of models that invoke full 3D slipping–reconnection and elongated ribbons (Pontin, Galsgaard, and Démoulin, 2016).

The good observing performance of the revamped Catania Solar Telescope has allowed us to investigate the evolution of the H α ribbon of the SOL2020-12-07T09:55 flaring event. Despite the low intensity of the flare, which was of B5.4 GOES class, the H α images showed the sensitivity of the Catania Solar Telescope to detect the variation of emission in time along the chromospheric ribbons, with a good reliability.

Indeed, the presence of the ribbon brightening in the chromosphere, which is also observed in the ultraviolet wavelengths, is one of the principal manifestations of the flare-energy release in the solar atmosphere. Using the Catania Solar Telescope and space-based instrument data we were also able to determine that this faint flare was characterized by the splitting of a single flux rope during the eruption, as usually observed in stronger events. Moreover, the temporal delay of the brightening observed in the chromosphere between




Figure 8 Sequence of IRIS SJIs at 1400, 1330, and 2796 Å acquired during the development of the SOL2020-12-07T09:55 flare, relative to the sub-FOV indicated with a dashed box in the $H\alpha$ map in Figure 5. All the images for the same passband are rescaled to the same minimum and maximum values.

the region along the PIL and the region corresponding to the northern footpoint of the filament allows us to infer the presence of a mechanism able to transfer the instability from the flux rope forming the main body of the filament to the northern footpoint of the filament.

Although the emission produced in the northern footpoint of the filament is only a few arcseconds away from the main ribbon, the reported delay of about 120 s between the peaks could be attributed to the twisted configuration of the magnetic field of the flux rope. In fact, the helical pattern of the flux rope, detected in the EUV images, could justify a longer time for the propagation of the signal in comparison to the typical time employed for the acceleration of the particles along a spine in a 3D null-point configuration (Romano et al., 2017).

This showcase clearly demonstrates the added value of the observations of the Catania Solar Telescope, which are useful both for space-weather forecasting and for scientific exploitation. In the near future, we plan to further upgrade the quality of the service provided by this telescope to the space-weather community by means of the implementation of techniques that improve the data quality, e.g., the lucky imaging technique, and an automatic detection algorithm for flare occurrences.

Acknowledgements This research received funding from the European Union's Horizon 2020 Research and Innovation 531 program under grant agreements No 824135 (SOLARNET) and support by the Italian Space Agency (ASI) under contract 2021-12-HH.0 to the co-financing INAF for the Italian contribution to the Solar-C EUVST preparatory science programme. This work was also supported by European Space Agency (ESA) in the context of the SWESNET project (contract 4000134036/21/D/MRP), by the Italian MIUR-PRIN 2017 on "Space Weather: impact on circumterrestrial environment of solar activity" and by the Università degli Studi di Catania (Piano per la Ricerca Università di Catania - Linea di intervento 2 "PIACERI").

Data Availability The datasets analyzed during the current study are available in the INAF-OACT repository at http://ssa.oact.inaf.it/oact/image_archive.php. For the flare description we also used IRIS data taken from <https://iris.lmsal.com/data.html>, HMI and AIA data available at <http://jsoc.stanford.edu/>.

Declarations

Disclosure of Potential Conflicts of Interest The authors declare that there are no conflicts of interest.

References

- Aulanier, G., Janvier, M., Schmieder, B.: 2012, *Astron. Astrophys.* **543**, A110. DOI.
- Aulanier, G., Démoulin, P., Schrijver, C.J., Janvier, M., Pariat, E., Schmieder, B.: 2013, *Astron. Astrophys.* **549**, A66. DOI.
- Barnes, G., Leka, K.D., Schrijver, C.J., Colak, T., Qahwaji, R., Ashamari, O.W., et al.: 2016, *Astrophys. J.* **829**, 89. DOI.
- Benz, A.O.: 2017, *Living Rev. Solar Phys.* **14**, 2. DOI.
- Bloomfield, D.S., Higgins, P.A., McAteer, R.T.J., Gallagher, P.T.: 2012, *Astrophys. J.* **747**, L41. DOI.
- De Groof, A., Walsh, A., Williams, D.: 2019, Metadata definition for Solar Orbiter science data. ESA. <https://issues.cosmos.esa.int/solarorbiterwiki/display/SOSP/Metadata+Definition+for+Solar+Orbiter+Science+Data>.
- De Pontieu, B., Title, A.M., Lemen, J.R., Kushner, G.D., Akin, D.J., Allard, B., et al.: 2014, *Solar Phys.* **289**, 2733. DOI.
- Falco, M., Costa, P., Romano, P.: 2019, *J. Space Weather Space Clim.* **9**, A22. DOI.
- Falconer, D.A., Moore, R.L., Gary, G.A.: 2008, *Astrophys. J.* **689**, 1433. DOI.
- Fletcher, L., Dennis, B.R., Hudson, H.S., Krucker, S., Phillips, K., Veronig, A., et al.: 2011, *Space Sci. Rev.* **159**, 19. DOI.
- Gibson, S.E., Fan, Y.: 2006, *Astrophys. J.* **637**, 65. DOI.
- Guglielmino, S.L., Zuccarello, F., Romano, P., Cristaldi, A., Ermolli, I., Criscuoli, S., et al.: 2016, *Astrophys. J.* **819**, 157. DOI.
- Kliem, B., Török, T., Titov, V.S., Lionello, R., Linker, J.A., Liu, R., et al.: 2014, *Astrophys. J.* **792**, 107. DOI.
- Kopp, R.A., Pneuman, G.W.: 1976, *Solar Phys.* **50**, 85. DOI.
- Korsós, M.B., Erdélyi, R.: 2016, *Astrophys. J.* **823**, 153. DOI.
- Korsós, M.B., Yang, S., Erdélyi, R.: 2019, *J. Space Weather Space Clim.* **9**, A6. DOI.

- Korsós, M.B., Ludmány, A., Erdélyi, R., Baranyi, T.: 2015, *Astrophys. J.* **802**, L21. DOI.
- Kuhn, K.R., Lin, H., Loran, D.: 1991, *Publ. Astron. Soc. Pac.* **103**, 1097. DOI.
- Law, N.M., Baldwin, J.E., Mackay, C.D.: 2006, *Astron. Astrophys.* **446**, 739. DOI.
- Leenaarts, J., Carlsson, M., Rouppe van der Voort, L.: 2012, *Astrophys. J.* **749**, 136. DOI.
- Lemen, J.R., Title, A.M., Akin, D.J., Boerner, P.F., Chou, C., Drake, J.F., et al.: 2012, *Solar Phys.* **275**, 17. DOI.
- Li, J.-W., Li, H., Li, Y., Feng, L., Huang, Y., Zhao, J., et al.: 2021, *Res. Astron. Astrophys.* **21**, 121. DOI.
- Mackay, C.D.: 2013, *Mon. Not. Roy. Astron. Soc.* **432**, 702. DOI.
- Masson, S., Pariat, E., Aulanier, G., Schrijver, C.J.: 2009, *Astrophys. J.* **700**, 559. DOI.
- Masson, S., Aulanier, E., Pariat, E., Klein, K.L.: 2012, *Solar Phys.* **276**, 199. DOI.
- Moore, R.L., Sterling, A.C., Hudson, H.S., Lemen, J.R.: 2001, *Astrophys. J.* **552**, 833. DOI.
- O'Dwyer, B., Del Zanna, G., Mason, H.E., Weber, M.A., Tripathi, D.: 2010, *Astron. Astrophys.* **521**, A21. DOI.
- Patsourakos, S., Georgoulis, M.K., Vourlidas, A., Nindos, A., Sarris, T., Anagnostopoulos, G., et al.: 2016, *Astrophys. J.* **817**, 14. DOI.
- Pesnell, W.D., Thompson, B.J., Chamberlin, P.C.: 2012, *Solar Phys.* **275**, 3. DOI.
- Piersanti, M., Alberti, T., Bemporad, A., Berrilli, F., Bruno, R., Capparelli, V., et al.: 2017, *Solar Phys.* **292**, 169. DOI.
- Pontin, D., Galsgaard, K., Démoulin, P.: 2016, *Solar Phys.* **291**, 1739. DOI.
- Romano, P., Elmhamdi, A., Kordi, A.S.: 2019, *Solar Phys.* **294**, 4. DOI.
- Romano, P., Zuccarello, F.: 2007, *Astron. Astrophys.* **474**, 633. DOI.
- Romano, P., Zuccarello, F., Guglielmino, S.L., Berrilli, F., Bruno, R., Carbone, V., et al.: 2015, *Astron. Astrophys.* **582**, A55. DOI.
- Romano, P., Falco, M., Guglielmino, S.L., Murabito, M.: 2017, *Astrophys. J.* **837**, 173. DOI.
- Romano, P., Elmhamdi, A., Falco, M., Costa, P., Kordi, A.S., Al-Trabulsi, H.A., et al.: 2018, *Astrophys. J.* **852**, L10. DOI.
- Scherrer, P.H., Schou, J., Bush, R.I., Kosovichev, A.G., Bogart, R.S., Hoeksema, J.T., et al.: 2012, *Solar Phys.* **275**, 207. DOI.
- Schwenn, R.: 2006, *Living Rev. Solar Phys.* **3**, 2. DOI.
- Ternullo, M., Contarino, L., Romano, P., Zuccarello, F.: 2006, *Astron. Nachr.* **327**, 36. DOI.
- Yuan, Y., Shih, F.Y., Jing, J., Wang, H.-M.: 2010, *Res. Astron. Astrophys.* **10**, 785. DOI.
- Zuccarello, F., Ermolli, I., Korsos, M., Giorgi, F., Guglielmino, S.L., Erdélyi, R., et al.: 2021, *Res. Astron. Astrophys.* **21**, 313. DOI.